


Modification of the jet properties at the Relativistic Heavy Ion Collider



Jan Rak

Department of Physics and Astronomy
Iowa State University


QCD in Heavy Ion collisions

Two particles azimuthal correlations in **pp**, **dAu** and **AuAu**

- nuclear modification of jet properties:
 - Intrinsic momentum k_T
 - Jet transverse fragmentation momentum j_T
 - Parton distribution function
 - Fragmentation function

Hard scattering

Hard scattering in transverse plane




Point-like partons \Rightarrow elastic scattering $\vec{p}_{T,jet1} + \vec{p}_{T,jet2} = \vec{0}$

Partons have intrinsic transverse momentum k_T $\vec{p}_{T,jet1} + \vec{p}_{T,jet2} = \vec{k}_{T,1} + \vec{k}_{T,2}$

Jet Fragmentation (width of the jet cone)

Partons have to materialize
(fragment) in colorless world

\vec{j}_T = jet fragmentation
transverse momentum



j_T and k_T are 2D vectors. We measure the mean value of its **projection** into the transverse plane $\langle |j_{Ty}| \rangle$ and $\langle |k_{Ty}| \rangle$.

$$\langle |k_{Ty}| \rangle = \sqrt{\frac{2}{\pi}} \sqrt{\langle k_T^2 \rangle}$$

$\langle |j_{Ty}| \rangle$ is an important jet parameter. It's constant value independent on fragment's p_T is characteristic of jet fragmentation (j_T -scaling).

$\langle |k_{Ty}| \rangle$ (intrinsic + NLO radiative corrections) carries the information on the parton interaction with QCD medium.

$$\langle k_{\perp}^2 \rangle_{AA} = \langle k_{\perp}^2 \rangle_{\text{vac}} + \langle k_{\perp}^2 \rangle_{\text{IS nucl}} + \langle k_{\perp}^2 \rangle_{\text{FS nucl}}$$


p+p

p+A

A+A

Fragmentation Function (distribution of parton momentum among fragments)

In Principle




$$\vec{p}_{parton} = \sum_i \vec{p}_i \quad |\vec{p}_{parton}| = \sum_i |\vec{p}_i| \cos(\theta_i)$$

$$z_i = \frac{|\vec{p}_i| \cos(\theta_i)}{|\vec{p}_{parton}|} \quad \sum_i z_i = 1$$

Fragmentation function $D(z) \propto e^{-z/\langle z \rangle}$

In Practice parton momenta are not known

$$x_E = -\frac{\vec{p}_T \cdot \vec{p}_{Ttrigg}}{|\vec{p}_{Ttrigg}|^2}$$



$$x_E z_{trigg} = \frac{p_T \cos(\Delta\phi)}{p_{parton}} = z$$

⇒ Simple relation


$$\langle z \rangle = \langle x_E \rangle \langle z_{trigg} \rangle$$

x_E in pp collisions


CCOR (ISR) $\sqrt{s} = 63$ GeV

see A.L.S. Angelis, Nucl Phys B209 (1982)

PHENIX preliminary



$1/\langle x_E \rangle \approx -4$ to -5



$1/\langle x_E \rangle \approx -5.3$

$\langle z \rangle$ extracted from pp data

We measured x_E and

$$\langle z \rangle = \langle x_E \rangle \langle z_{trigg} \rangle$$

$$x_{Ttrigg} = 2 \cdot p_{Ttrigg} / \sqrt{s}$$

$$\langle z_{trigg} \rangle \propto \int_{x_{Ttrigg}}^1 z \cdot e^{-z/\langle z \rangle} f_q(p_T / z) \cdot z^{-2} dz$$


Only one unknown variable $\langle z \rangle \Rightarrow$ iterative solution

FFn $D(z)$

extracted from PHENIX



$p+p \rightarrow \pi^0 + X$

Slope of the fragmentation function
in p+p collisions at $\sqrt{s}=200$ GeV



$$\frac{1}{\langle z \rangle} = 6.16 \pm 0.32$$

pp and dAu correlation functions





Fixed correlation:

both $p_{T\text{trigg}}$ and $p_{T\text{assoc}}$ are in the same range

Assorted correlation:

$p_{T\text{trigg}}$ and $p_{T\text{assoc}}$ different



5.0 < $p_{T\text{trigg}}$ < 16.0 GeV/c

Jet function assumed to be Gaussian

$$C_{ij}(\Delta\phi) = \text{norm} \cdot \frac{dN_{ij}^{\text{real}}}{d\Delta\phi_{ij}} / \frac{dN_{ij}^{\text{mixed}}}{d\Delta\phi_{ij}}$$



$$\text{Fit} = \text{const} + \text{Gauss}(0) + \text{Gauss}(\pm\pi)$$

$\sigma_N, \sigma_A, \langle |j_{Ty}| \rangle, \langle |k_{Ty}| \rangle$ relations

Knowing σ_N and σ_A it is straightforward to extract $\langle |j_{Ty}| \rangle$ and $\langle z_{trigg} \rangle \langle |k_{Ty}| \rangle$

In the high- p_T limit ($p_T \gg \langle |j_{Ty}| \rangle$ and $p_T \gg \langle |k_{Ty}| \rangle$)

$$\langle |j_{\perp y}| \rangle = \langle p_{\perp} \rangle \sin \frac{\sigma_N}{\sqrt{\pi}}$$


$$\langle |k_{Ty}| \rangle \approx \langle p_T \rangle \sqrt{\sigma_A^2 - \sigma_N^2}$$

However, inspired by Feynman, Field, Fox and Tannenbaum (see *Phys. Lett. 97B (1980) 163*) we derived more accurate equation

$$\langle z_{trigg} \rangle \langle |k_{Ty}| \rangle = \frac{\langle p_T \rangle}{\sqrt{2} x_h} \sqrt{\sin^2 \sqrt{\frac{2}{\pi}} \sigma_A - (1 + x_h^2) \sin^2 \frac{\sigma_N}{\sqrt{\pi}}}$$

$$x_h = p_{T,assoc} / p_{T,trigg}$$

$\sigma_N, \sigma_A \rightarrow \langle |j_{Ty}| \rangle, \langle |k_{Ty}| \rangle$ in pp data



$$\langle k_{\perp}^2 \rangle_{pp} = \langle k_{\perp}^2 \rangle_{vac}$$




PHENIX preliminary

$$\langle |j_{Ty}| \rangle = 359 \pm 11 \text{ MeV/c}$$


$$\langle |k_{Ty}| \rangle = 964 \pm 49 \text{ MeV/c}$$

Both $\langle |j_{Ty}| \rangle$ and $\langle |k_{Ty}| \rangle$ in very good agreement with previous measurements:
PLB97 (1980)163
PRD 59 (1999) 074007


Di-jet fragmentation



"wave function collapse"



$$f_q(p_{Tq}, p_{Tassoc}) = D\left(\frac{p_{Tassoc}}{p_{Tq}}\right) \cdot p_{Tq}^{-8}$$



For fixed p_T correlation the parton distribution function is almost unaffected by the condition of having fragment on the opposite side.

kT-smearing

Associated parton distribution $f_a = kT \otimes f_t$ and the final formula for invariant cross section is

$$\frac{1}{p_t} \frac{d^2\sigma}{dp_t dp_a} = \frac{1}{p_t} \int_{p_t}^{\sqrt{s}/2} dq_t \cdot D\left(\frac{p_t}{q_t}\right) \cdot \frac{1}{q_t} \int_{p_a}^{\sqrt{s}/2} dq_a \cdot C_{bb}(q_a - q_t, \langle k_T^2 \rangle) \cdot f(q_a) \cdot D\left(\frac{p_a}{q_a}\right)$$

$$q_t \rightarrow \frac{p_t}{z_t}$$

Correlation fcn
between b2b partons
gaussian for $k_T > 0$ or delta
function for $k_T = 0$

$$\frac{1}{p_t} \frac{d^2\sigma}{dp_t dp_a} = \frac{1}{p_t} \int_{p_t}^{\sqrt{s}/2} dz_t \cdot D(z_t) \cdot \frac{1}{z_t} \int_{p_a}^{\sqrt{s}/2} dq_a \cdot C_{bb}\left(q_a - \frac{p_t}{z_t}, \langle k_T^2 \rangle\right) \cdot f(q_a) \cdot D\left(\frac{p_a}{q_a}\right)$$



Di-jet mean z

conditional

$$\langle z_t \rangle = \frac{\int_{x_{Tt}}^{p_{Tt}/p_{Ta}} f_q\left(\frac{p_{Tt}}{z_t}\right) \cdot D(z_t) \cdot D\left(\frac{p_{Ta}}{p_{Tt}} z_t\right) \cdot dz_t}{\int_{x_{Tt}}^{p_{Tt}/p_{Ta}} \frac{1}{z_t} f_q\left(\frac{p_{Tt}}{z_t}\right) \cdot D(z_t) \cdot D\left(\frac{p_{Ta}}{p_{Tt}} z_t\right) \cdot dz_t}$$


inclusive

$$\langle z \rangle = \frac{\int_{x_{\perp}}^1 z z^{-2} D(z) f_q(p_{\perp}/z) dz}{\int_{x_{\perp}}^1 z^{-2} D(z) f_q(p_{\perp}/z) dz}$$





Simulation

Sorry, I can show only simulation – the final data Phys. Rev. C.




PHENIX/CCOR p_{trigg} slopes




Associated yields - complementary method of $\langle |k_{Ty}| \rangle$ extraction

PHENIX $\langle |k_{Ty}| \rangle = 0.9$ GeV/c



$\frac{1}{N} \frac{dn_2}{dp_a}$



STAR $\langle |k_{Ty}| \rangle = 1.2$ GeV/c


We can describe all p_{Tassoc} distribution with unique value of $\langle |k_{Ty}| \rangle = 0.9$ GeV/c, but not STAR data.

AuAu $\langle |j_{Ty}| \rangle$ and $\langle z \rangle \langle |k_{Ty}| \rangle$ from CF

Jana Bielcikova

Phys.Rev.Lett.92:032301,2004



$$\langle k_{\perp}^2 \rangle_{AA} = \langle k_{\perp}^2 \rangle_{vac} + \langle k_{\perp}^2 \rangle_{IS\ nucl} + \langle k_{\perp}^2 \rangle_{FS\ nucl}$$



There seems to be significant broadening of the away-side correlation peak which persists also at somewhat higher p_T range.

AuAu associated yields

$$(2.5 < p_{T\text{trigg}} < 4.0) \otimes (1.0 < p_{T\text{assoc}} < 2.5) \text{ GeV/c}$$




Note p_T is rather low; associated particle yields increase with centrality

p_T distributions on near and away side

Near side:

**Overall
enhancement
from pp to AA**



Away side:

energy from
initial parton seems
to be converted
to lower p_T particles

reminiscent of energy
loss predictions

Apparent modification of the fragmentation function ?

Summary and conclusions

Jet production and fragmentation:

- Good agreement of the jet properties in pp collisions with other lower \sqrt{s} experiments
- dAu j_T and k_T consistent with pp
- In AuAu significant broadening of “effective” k_T - with centrality
- Yield of away side associated particles is suppressed at $p_T > 2\text{GeV}/c$ and shows rising trend with N_{part} below $2\text{GeV}/c$. Remnant of high- p_T jets - hint of jet-quenching balance ?

